

The projective class ring of half-quantum groups at roots of unity

Claude Cibils

Abstract

We describe the structure of the Grothendieck ring of projective modules of basic Hopf algebras using a positive integer determined by the composition series of the principal indecomposable projective module.

1991 Mathematics Subject Classification : 16E20 and 16W30

1 Introduction

We consider finite dimensional Hopf algebras which are basic and split, which means that simple module are one-dimensional over the ground field. Some of those algebras have been called half-quantum groups and are useful in knot theory and 3-manifold invariants, see [7, 13]. Through the Drinfeld quantum double ([9]) they provide universal R -matrices and solutions to the Yang-Baxter equation. Notice that the relation between the K-theory of a Hopf algebra and the K-theory of its quantum double is not understood. Basic and split Hopf algebras has been studied in [6, 5, 10].

The purpose of this paper is to describe the projective class ring of an arbitrary basic and split finite dimensional Hopf algebra, where the multiplicative structure is given by the tensor product of modules obtained through the comultiplication of the Hopf algebra. We obtain for instance that for q a root of unity, the Grothendieck ring of $u_q^+(sl_2)$ has nilpotent elements. Notice that $u_q^+(sl_2)$ has only a finite number of isomorphism classes of indecomposable modules; by contrast Benson and Carlson proved in [2] that a group algebra verifying this property has no nilpotent elements in its Grothendieck ring.

The tensor product of simple modules is often commutative, this happens to be true for half-quantum groups at roots of unity. With this assumption,

the structure of the complexification of the projective class ring can be made more precise than in the general case, see Theorem 3.12. The result relies on a positive integer determined by the number of roots of unity where the composition series of the projective cover of the trivial module do not vanish. Its value for sl_2 is 1, and the complexification of the projective class ring of sl_2 is the product of two copies of the complex numbers with the product of $n - 1$ copies of the dual numbers.

The projective class ring of half-quantum groups at roots of unity are commutative rings. Actually the entire Grothendieck ring of $u_q^+(sl_2)$ is commutative, as a consequence of direct computations made in [4]; there is no theoretical result predicting this commutativity property since it is known that half-quantum groups are not quasitriangular, and not even quasi-cocommutative (see [14]).

2 Basic and split Hopf algebras

Let Λ be a finite dimensional Hopf algebra over a field k , that is an associative algebra, provided with a comultiplication $\Delta : \Lambda \rightarrow \Lambda \otimes \Lambda$, a counit $\varepsilon : \Lambda \rightarrow k$ and an antipode $S : \Lambda \rightarrow \Lambda$ (see [16]).

The tensor product $M \otimes N$ over k of two left Λ -modules M and N has a left Λ -module structure obtained through Δ . The ground field k_ε equipped with the module structure given by ε is the unit element; since $M^* = \text{Hom}_k(M, k)$ is a right Λ -module, it turns into a left module by composing the action with the antipode.

An associative algebra is split if the endomorphism rings of the simple modules are isomorphic to the ground field; this condition is always satisfied if k is algebraically closed. The algebra is basic if each simple module has multiplicity one in its maximal semisimple quotient algebra. Each basic and split Hopf algebra has a structure group attached, which corresponds to the group of group-likes in the dual Hopf algebra.

Lemma 2.1 *Let $\mathcal{S}$ be the set of isomorphism classes of simple modules over a basic and split finite dimensional Hopf algebra. Then $\mathcal{S}$ is a group.*

Proof: Let S and T be simple modules. Since they are one-dimensional, their tensor product is also one-dimensional, hence a simple module. The unit element in $\mathcal{S}$ is the isomorphism class of k_ε , while the inverse of S is S^* .

Example 2.2 Let G be a finite group and k^G be the Hopf algebra of functions on it, that is the dual of the group algebra kG . Its structure group is G .

Example 2.3 Let $u_q^+(\mathcal{G})$ be the half-quantum group corresponding to a simple Lie algebra $\mathcal{G}$ with Cartan matrix of size $t \times t$, and to a primitive root of unity of order n . This basic and split algebra has an abelian structure group, isomorphic to the product of t copies of the cyclic group of order n .

If the algebra is not basic, the tensor product of simple modules is not semisimple in general. The structure group gives an easy proof of a particular case of a result due to P. Schauenburg. Recall that associative algebras are Morita equivalent if their module categories are equivalent. Hopf algebras are Morita-monoidally equivalent if their module categories are equivalent as monoidal categories, which means that there exist an equivalence of categories compatible with tensor products, dual operations and the unit object. In [15] Schauenburg shows that finite dimensional Hopf algebras Morita-monoidally equivalent have isomorphic underlying associative algebras.

Proposition 2.4 *Let Λ be a basic and split Hopf algebra, and let Λ' be a Hopf algebra Morita-monoidally equivalent to Λ . Then Λ and Λ' are isomorphic associative algebras.*

Proof: Consider the structure group $\mathcal{S}$ of Λ and the set $\mathcal{S}'$ of isomorphism classes of simple Λ' -modules. It follows from the definition that $\mathcal{S}'$ is also a group with respect to the tensor product of modules, isomorphic to $\mathcal{S}$. Since the dimension over the ground field is multiplicative on tensor products and since each simple module has an inverse, simple Λ' -modules are one-dimensional. Hence Λ' is basic and split algebra, and consequently isomorphic to Λ .

We recall now definitions which will be useful at the next section. The Grothendieck group of an abelian category $\mathcal{C}$ is the free abelian group with basis the isomorphism classes $[X]$ of objects X of $\mathcal{C}$, divided by the subgroup generated by elements $[X_2] - [X_1] - [X_3]$ provided by each split exact sequence $0 \rightarrow X_1 \rightarrow X_2 \rightarrow X_3 \rightarrow 0$ in $\mathcal{C}$. We denote the quotient group $K_0(\mathcal{C})$, and $K_0(\Lambda)$ denotes the Grothendieck group of the category of finitely generated modules over a ring Λ . In case $\mathcal{C}$ is a monoidal abelian category – for instance

the category of modules over a Hopf algebra – the free abelian group above is a ring. Tensoring a split exact sequence yields to a split exact sequence, the quoted subgroup is a two-sided ideal and the Grothendieck group is a ring.

Remark 2.5 If Λ is a finite dimensional algebra, $K_0(\Lambda)$ is a free abelian group with a basis given by the isomorphism classes of indecomposable modules, as an immediate consequence of the Krull-Schmidt Theorem (see for instance [8]).

Consider the subgroup generated by elements given by non necessarily split exact sequences and denote the corresponding quotient group $\overline{K}_0(\mathcal{C})$. For modules over a finite dimensional algebra Λ , the Jordan-Holder theorem insures that $\overline{K}_0(\Lambda)$ is free abelian with a basis provided by the isomorphism classes of simple modules. Since Λ is a Hopf algebra over a field k , the functors $X \otimes -$ and $- \otimes X$ are exact for any Λ module X , and $\overline{K}_0(\Lambda)$ is a ring. If Λ is a basic and split Hopf algebra, this ring is clearly isomorphic to the group ring $\mathbf{Z}\mathcal{S}$, where $\mathcal{S}$ is the structure group of Λ .

There is a canonical ring surjection $K_0(\Lambda) \rightarrow \overline{K}_0(\Lambda)$ called dimension vector which associates to a module M the sum $\overline{M}$ of the quotients of one of its composition series. This map is an isomorphism if and only if Λ is a semisimple algebra.

3 Projective class rings

We study $K_0^p(\Lambda)$, the subring of $K_0(\Lambda)$ generated by the projective and semisimple modules. We recall the following standard results:

Lemma 3.1 *Let Λ be a Hopf algebra, and let P and X be finitely generated modules such that P is projective. Then $P \otimes X$ is a projective module.*

Proof: We prove that $\Lambda \otimes X$ is free. The module $\Lambda \otimes X$ is isomorphic to $\Lambda \otimes X_\varepsilon$, where X_ε is the vector space X equipped with the trivial action given by the counit ε . The result follows since $\Lambda \otimes X_\varepsilon$ is free, of rank the dimension of X . An isomorphism of left Λ -modules $\varphi : \Lambda \otimes X \rightarrow \Lambda \otimes X_\varepsilon$ is given by $\varphi(a \otimes x) = \sum a_{(1)} \otimes S(a_{(2)})x$, where $\Delta a = \sum a_{(1)} \otimes a_{(2)}$.

Proposition 3.2 *Let Λ be a finite dimensional basic and split Hopf algebra. The subring $K_0^p(\Lambda)$ generated by semisimple and projective modules is a free abelian group with basis the set of isomorphism classes of simple and indecomposable projective modules.*

Proof: $K_0(\Lambda)$ is free abelian with basis the set of isomorphism classes of indecomposable modules. The described subgroup is the smallest containing semisimple and projective modules. Moreover this subgroup is an ideal, due to Lemma 3.1 and to the fact that isomorphism classes of simple modules forms a group.

Let $\mathcal{P}$ be the set of isomorphism classes of indecomposable projective modules. The ring $\overline{K}_0(\Lambda)$ is the integral group ring of the structure group $\mathcal{S}$, and $\mathbf{Z}\mathcal{P}$ is a $\mathbf{Z}\mathcal{S}$ -bimodule. It is well known that $\mathcal{S}$ and $\mathcal{P}$ are canonically bijective sets (see for instance [8]) : each simple module S has an unique projective cover P_S , a projective module such that there exist an essential surjection $\psi : P_S \rightarrow S$: for any strict submodule X of P_S , the restriction of ψ to X is not surjective.

Lemma 3.3 *For a basic and split finite dimensional Hopf algebra, $\mathbf{Z}\mathcal{P}$ is isomorphic to $\mathbf{Z}\mathcal{S}$ as a $\mathbf{Z}\mathcal{S}$ -bimodule.*

Proof: Consider the above canonical bijection : we have to prove that $T \otimes P_S$ is isomorphic to $P_{T \otimes S}$ for T and S simple modules. Since $\psi : P_S \rightarrow S$ is an essential surjection, we obtain that $T \otimes P_S \rightarrow T \otimes S$ is also essential since simple modules are one-dimensional. Moreover, $T \otimes P_S$ is a projective module, and the unicity of projective covers insures that $T \otimes P_S$ is the projective cover of $T \otimes S$.

Lemma 3.4 *Let Λ be a finite dimensional basic and split Hopf algebra which is not semisimple. Then $K_0^p(\Lambda)$ is the direct sum of the subring $\mathbf{Z}\mathcal{S}$ and the $\mathbf{Z}\mathcal{S}$ -bimodule $\mathbf{Z}\mathcal{P}$.*

Proof: Since $\overline{K}_0(\Lambda)$ is the subring $\mathbf{Z}\mathcal{S}$, we need to prove that $\mathcal{S} \cap \mathcal{P}$ is empty; indeed if there exist a simple and projective module, we infer that any simple module is projective by Lemma 3.1 and the fact that $\mathcal{S}$ is a group. If all the simple modules are projective modules, there are no non trivial extensions between them, and the algebra is semisimple.

Remark 3.5 A semisimple basic and split Hopf algebras is of the form k^G for G a finite group. Such algebras are commutative and $K_0^p(\Lambda)$ is the group algebra $\mathbf{Z}G$ since the structure group of k^G is G .

The next purpose is to describe the multiplicative structure of $\mathbf{Z}\mathcal{P}$, in other words to describe the $\mathbf{Z}\mathcal{S}$ -bimodule map

$$\mathbf{Z}\mathcal{P} \otimes_{\mathbf{Z}\mathcal{S}} \mathbf{Z}\mathcal{P} \longrightarrow \mathbf{Z}\mathcal{P}$$

induced by the tensor product of modules over a basic and split Hopf algebra.

Proposition 3.6 *Let Λ be any finite dimensional Hopf algebra, and let M and P be finitely generated left Λ -modules where P is projective. Then $M \otimes P$ is isomorphic to $\overline{M} \otimes P$.*

Proof: Recall that $\overline{M}$ is the semisimple module obtained by adding the quotients of a composition series of M . It suffices to prove that for an exact sequence

$$0 \rightarrow M_1 \rightarrow M \rightarrow M_2 \rightarrow 0$$

of Λ -modules we have that $M \otimes P$ is isomorphic to $M_1 \otimes P \oplus M_2 \otimes P$. By tensoring, we get an exact sequence

$$0 \rightarrow M_1 \otimes P \rightarrow M \otimes P \rightarrow M_2 \otimes P \rightarrow 0$$

of Λ -modules. Since $M_2 \otimes P$ is projective by Lemma 3.1, the sequence splits.

Corollary 3.7 *If the structure group of a basic and split finite dimensional Hopf algebra is abelian, the projective modules are central in $K_0(\Lambda)$.*

Proof: We prove that any indecomposable projective module is in the center of $K_0(\Lambda)$, more precisely we have that $M \otimes P_S$ and $P_S \otimes M$ are isomorphic modules for any finitely generated module M and any simple module S . From the proof of the preceding result we have that $P_S \otimes M$ and $P_S \otimes \overline{M}$ are isomorphic modules. We need to prove that $P_S \otimes T$ and $T \otimes P_S$ are isomorphic for any simple module T . The module $P_S \otimes T$ is the projective cover of $S \otimes T$ (see the proof of Lemma 3.3) and $T \otimes P_S$ the projective cover of $T \otimes S$. Since $\mathcal{S}$ is commutative, and projective covers are unique up to isomorphism, we obtain the result.

Another way of formulating Proposition 3.6 is to consider the Grothendieck group $K_0(\mathcal{P}\Lambda)$ of the category of projective finitely generated Λ -modules. The natural two-sided action of $K_0(\Lambda)$ on $K_0(\mathcal{P}\Lambda)$ provides an action of $\overline{K}_0(\Lambda)$ on $K_0(\mathcal{P}\Lambda)$.

Definition. Let A be a ring; a multiplicative A -bimodule is an A -bimodule M provided with an associative A -bimodule map $\varphi : M \otimes_A M \rightarrow M$.

If M is a multiplicative A -bimodule, then $A \oplus M$ has a natural ring structure where M is a two-sided ideal. Conversely, if R is a ring with a

subring A and a decomposition $R = A \oplus I$ where I is a two-sided ideal of R , we obtain that I is a multiplicative A -bimodule. The ring $K_0^p(\Lambda)$ corresponds to this situation, since $K_0^p(\Lambda) = \mathbf{Z}\mathcal{S} \oplus \mathbf{Z}\mathcal{P}$. Moreover, $\mathbf{Z}\mathcal{P}$ is a $\mathbf{Z}\mathcal{S}$ -bimodule isomorphic to $\mathbf{Z}\mathcal{S}$; we need to study the multiplicative bimodule structures of a ring having underlying bimodule the ring itself.

Proposition 3.8 *The multiplicative bimodule structures on a ring A are in bijection with its center ZA . The isomorphism classes of bimodule structures on A are in one-to-one correspondance with $ZA/U(ZA)$, where $U(ZA)$ is the group of units of A .*

Proof: Notice that an A -bimodule map $\varphi : A \otimes_A A \rightarrow A$ is an A -bimodule endomorphism of A which is completely determined by the image of 1, an element of the center of A . The associativity always holds since

$$\varphi(\varphi(a \otimes b) \otimes c) = abc \varphi(1)^2 = \varphi(a \otimes \varphi(b \otimes c)).$$

Finally an A -bimodule automorphism of A is given by a unit of ZA , we infer that two multiplicative bimodule structures φ and φ' are isomorphic if and only if there exist a unit $u \in U(ZA)$ such that $\varphi(1) = u\varphi'(1)$.

We define now a canonical element c of the integral group ring $\mathbf{Z}\mathcal{S}$ as follows: let P_1 be the projective cover of the trivial module k_ϵ (according to the preceding notation we have $P_1 = P_{k_\epsilon}$). The element c is provided by a composition series of P_1 , more precisely $c = \overline{P_1}$. Notice that for each $S \in \mathcal{S}$, we have $\overline{P_S} = Sc$ in the group ring.

Proposition 3.9 *Let Λ be a basic and split Hopf algebra, with structure group $\mathcal{S}$ (non necessarily abelian) and canonical element c . Then $K_0^p(\Lambda)$ is a ring isomorphic to $\mathbf{Z}\mathcal{S} \oplus \mathbf{Z}\mathcal{S}$, where the first factor is the group ring and multiplication is given by :*

$$(s_1, t_1)(s_2, t_2) = (s_1s_2, s_1t_2 + t_1s_2 + t_1ct_2).$$

Example 3.10 Let q be a primitive n -th root of unity in $\mathbf{C}$, and let $u_q^+(sl_2)$ be the corresponding finite dimensional half-quantum group, which structure group $\mathcal{S}$ is isomorphic to the cyclic group of order n (see [6]). The element c is the trace element in $\mathbf{Z}\mathcal{S}$, i.e. the sum of all the group elements (the structure of $u_q^+(sl_2)$ -modules is given in [4]).

Corollary 3.11 *The canonical element c of a basic and split non semisimple Hopf algebra lies in the center of the integral group ring of its structure group.*

In case the structure group is commutative there is a more explicit description of the algebra $\mathbf{C} \otimes K_0^p(\Lambda)$. Recall that $\mathbf{CS}$ is a semisimple commutative algebra isomorphic to $\mathbf{C}^{\mathcal{S}} = \times_{S \in \mathcal{S}} \delta_S$, where $\{\delta_S\}_{S \in \mathcal{S}}$ is the complete set of primitive orthogonal idempotents provided by the Dirac masses on elements of $\mathcal{S}$. An explicit isomorphism is provided by the Fourier transform once generators of $\mathcal{S}$ and corresponding roots of unity are given.

Definition: Let Λ be a basic and split Hopf algebra with abelian structure group $\mathcal{S}$ and canonical element c . The positive integer r is the number of non-zero coordinates of c in the above decomposition of the group algebra $\mathbf{CS}$. Equivalently, r is the number of irreducible characters of $\mathcal{S}$ which do not vanish at c .

Theorem 3.12 *Let Λ be a basic and split Hopf algebra with abelian structure group of order s . Then $\mathbf{C} \otimes K_0^p(\Lambda)$ is isomorphic to the algebra $\mathbf{C}^{2r} \times \mathbf{C}[\epsilon]^{s-r}$ where $\mathbf{C}[\epsilon]$ is the algebra of dual numbers $\mathbf{C}[X]/X^2$.*

Proof: We know that a multiplicative bimodule structure on a ring A having underlying bimodule A is determined by a central element up to multiplication by a unit of the center. After normalization we have to describe the algebra structure of $\mathbf{C}^s \oplus \mathbf{C}^s$ where the first factor affords its natural algebra structure and the second is considered as a $\mathbf{C}^s$ -bimodule with multiplicative structure provided by an element c' having r entries with value 1, the others entries having zero value. For simplicity we consider $\mathbf{C}^s$ as the algebra of functions $\mathbf{C}^E$ on the set E , and c' be given by a subset F in E . A complete set of primitive orthogonal idempotents is

$$\{(\delta_y, 0)\}_{y \in E \setminus F} \sqcup \{(\delta_x, -\delta_x), (0, \delta_x)\}_{x \in F}.$$

Moreover, the elements $\{(0, \delta_y)\}_{y \in E \setminus F}$ are orthogonal elements with zero square. Actually

$$(\delta_y, 0)(0, \delta_y) = (0, \delta_y) \text{ if } y \in E \setminus F$$

$$\text{while } (\delta_x, -\delta_x)(0, \delta_y) = 0 = (0, \delta_x)(0, \delta_y) \text{ for } y \in E \setminus F \text{ and } x \in F.$$

This formulas are equivalent to the structure result for $K_0^p(\Lambda)$.

Remark 3.13 For $u_q^+(sl_2)$, we notice that c is the sum of all the elements of the cyclic group of order n , hence $r = 1$. Then

$$\mathbf{C} \otimes K_0^p(u_q^+(sl_2)) \cong \mathbf{C}^2 \times \mathbf{C}[\epsilon]^{n-1}.$$

Finally we describe the virtual projective modules which have zero square. Consider G a finite abelian group with a given system of generators

$$G = \langle K_1, \dots, K_t \mid K_i^{n_i} = 0, K_i K_j = K_j K_i \rangle,$$

and let $q_1, \dots, q_t$ be complex primitive roots of unity of order $n_1, \dots, n_t$. If $a = (a_1, \dots, a_t)$ is a t -uple of integers, we put $q^a = q_1^{a_1} \cdots q_t^{a_t}$, and $K^a = K_1^{a_1} \cdots K_t^{a_t}$. Moreover $ab = \sum_{i=1}^t a_i b_i$ for $b = (b_1 \cdots b_t)$.

Consider the non-degenerate bilinear form

$$\beta_G : \mathbf{C}G \times \mathbf{C}G \longrightarrow \mathbf{C}$$

given by the isomorphism $\alpha : \mathbf{C}G \rightarrow \mathbf{C}^G$ of Hopf algebras, namely

$$\beta_G(K^a, K^b) = q^{ab}.$$

Let Λ be a basic and split Hopf algebra with structure group $\mathcal{S}$ and canonical element c . The ideal of projective modules in $\mathbf{C} \otimes K_0^p(\Lambda)$ is isomorphic to $(\mathbf{C}\mathcal{S})_c$ where the product is given by $t_1.t_2 = t_1 c t_2$. Let $B_c = \{K^x \in \mathcal{S} \mid \beta_{\mathcal{S}}(K^x, c) \neq 0\}$.

Proposition 3.14 *The nilradical of $\mathbf{C} \otimes K_0^p(\Lambda)$ is the orthogonal vector space to $\mathbf{C}B_c$ in $\mathbf{C}\mathcal{S}$.*

Proof : Consider the corresponding product in $\mathbf{C}^{\mathcal{S}}$, given by $f.g = f\alpha(c)g$. An element f has zero square if $0 = f.f = ff\alpha(c)$, which only holds for $f\alpha(c) = 0$. This means that for $K^x \in B_c$, we have $f(K^x) = 0$. The corresponding element in $\mathbf{C}\mathcal{S}$ is orthogonal to each element of B_c .

Example 3.15 For $u_q^+(sl_2)$ the element c is the sum of all the elements of the cyclic group of order n , then $B_c = \{1\}$. The orthogonal space is the augmentation ideal and any difference of two indecomposable projective modules has zero square.

References

- [1] Anderson, F.W., Fuller, K.R.: Rings and categories of modules, Springer GTM 13, New York 1973
- [2] Benson, D.J., Carlson, J.F. Nilpotent elements in the Green ring. *J. Algebra* **104**, 329–350 (1986)
- [3] Chari, V., Pressley, A.: A guide to quantum groups. Cambridge University Press. 1994
- [4] Cibils, C.: A quiver quantum group. *Commun. in Math. Phys.* **157**, 459–477 (1993)
- [5] Cibils, C., Rosso, M.: Algèbres des chemins quantiques. *Advances in Math.* **125**, 171–199 (1997)
- [6] Cibils, C.: Half-quantum groups at roots of unity, path algebras and representation type. *International Mathematics Research Notices* **1997:12**, 541–553 (1997)
- [7] Crane, L., Frenkel, I.B.: Four-dimensional topological quantum field theory, Hopf categories, and the canonical bases. *J. Math. Phys.* **35**, 5136–5154 (1994)
- [8] Curtis, C.W., Reiner, I.: Methods of representation theory, Pure & applied mathematics, Wiley-interscience, New York, 1981
- [9] Drinfel'd, V.G.: Quantum groups. *Proceedings of the International Congress of Mathematicians*. Vol. 1, pp. 798–820. Berkeley: Academic Press 1986
- [10] Green, E., Solberg, O. Basic Hopf algebras and quantum groups, to appear in *Math. Zeit.*
- [11] Green, E.N., Marcos, E., Solberg, O.: Representations and almost split sequences for Hopf Algebras, *Representation theory of algebras*, R. Bautista, R. Martinez-Villa, J. de la Peña, Editors, *Cand. Math. Soc., Conference Proceedings*, **18** 237–245 (1996)
- [12] Kassel, C.: Quantum groups. *Grad. Texts in Math.* **155** 1995
- [13] Kuperberg, G. Non-involutory Hopf algebras and 3-manifold invariants. *Duke Math J.* **84**, 83–129 (1996)
- [14] Radford, D.E.: Minimal quasitriangular Hopf algebras. *J. Algebra* **157**, 285–315 (1993)
- [15] Schauenburg, P.: Hopf bigalois extensions. *Communications in Algebra* **24**, 3797–3825 (1996)

[16] Sweedler, M.E.: Hopf algebras. Benjamin. New York U.S.A. 1969

Dép. de Mathématiques, Université de Montpellier 2, F-34095 Montpellier cedex 5.
`cibils@math.univ-montp2.fr`

February 1998